

Homosexuality

(When quoting scriptures, from the Rotherham Emphasized Bible New Testament, I will substitute the Hebrew words Yahoshua (yeh-ho-shoo'- ah) for Jesus, Yahweh and Elohim for God and the LORD and ruah for pneuma (spirit).)

The great deceiver has dressed up, groomed and perfumed the lifestyle of homosexuality, presenting it as an alternative, gay, free lifestyle when it is actually death and total bondage, shackling all whom it seduces. Yahweh, who ahabs (loves) man, who is an Elohim of Life, has warned man, from the beginning of time, to flee from sexual immorality. Today, homosexuality presents a grave threat in many cultures. It is being introduced to our youths and encouraged by the President of the United States, Barack Hussein Obama, our schools, television and actors, to name only a few of its avid supporters. Yahweh's chief enemy, the god of this age is behind this spiritual assault, which must be countered by the truth of Yahweh's Word.¹ Truths from the Word of Yahweh give us Life, setting us free while man's deceptions bind, maim and destroy those who give heed. If the Word of Yahweh is not enough proof that homosexuality is death, the medical field will also confirm this absolute universal truth.

In our day, the consequences of disobeying Yahweh by accepting homosexuality has murdered millions. "More than 18,000 people with AIDS still die each year in the US. Gay, bisexual, and other men who have sex with men (MSM)[†] are strongly affected and represent the majority of persons who have died. Through 2007, more than 576,000 people with AIDS in the US have died since the epidemic began."²

AIDS Diagnoses by Age in the United States³

Age (Years)	Estimated # of AIDS Diagnoses, 2009	Cumulative Estimated # of AIDS Diagnoses, through 2009*
Under 13	13	9,448
Ages 13-14	58	1,321
Ages 15-19	484	7,214
Ages 20-24	2,095	42,920
Ages 25-29	3,476	129,639
Ages 30-34	4,043	214,149
Ages 35-39	4,893	234,575
Ages 40-44	5,689	193,237
Ages 45-49	5,466	126,380
Ages 50-54	3,983	72,327
Ages 55-59	2,191	39,025
Ages 60-64	1,010	20,633
Ages 65 or older	846	17,743

¹ Eph. 6:10-18

² <http://www.cdc.gov/hiv/resources/factsheets/us.htm>

³ <http://www.cdc.gov/hiv/topics/surveillance/basic.htm#aidsage>

In the United States, the NEA, the union for public school teachers, has now embraced homosexuality.⁴ Their stealth technique of accepting and teaching this wickedness into our public schools is by treating homosexuals as a racial group who are being harassed and bullied for their chosen lifestyle. These teachers condemn the smoking of cigarettes but are silent on the consequences of homosexual behavior, such as the extraordinarily high disease, suicide and death rate, as was mentioned above. They are teaching the children that these lifestyles are now accepted and may even be encouraged; by doing so, these teachers are leading the innocent sheep to the butchers (pedophiles).⁵ ("Although most homosexual activists publicly deny that they want access to boys, many homosexual groups around the world are working aggressively to lower the age of sexual consent. Their cause is being aided by the professional psychiatric and psychological associations, which have moved in recent years toward normalizing pedophilia, much as they did with homosexuality in the early 1970s."⁶)

Homosexual means, "Involving, related to, or characterized by a sexual propensity for one's own sex; of or involving sexual activity with a member of one's own sex, or between individuals of the same sex."⁷ Homosexuality is nothing new but the names have been changed to recruit new members; names such as 'gay,' 'lesbian' and 'transgender' reduce the negative image homosexuality has with it. Yahweh deals with it, in the book of Genesis, by destroying the inhabitants of Sodom and Gomorrah, who were described as being excellently base and sinful (Gen. 19).⁸ The word, 'sodomy,' comes from the wickedness that was transpiring at Sodom. Lot had two messengers (angels) visit him when the men of Sodom demanded of Lot, "...Bring them forth unto us, and let us know them" (Gen. 19:5). Yahweh had Moses write about sexual immorality:

A woman shall not have on, the wearing apparel of a man, nor shall a man put on the mantle of a woman; for, an abomination to Yahweh thy Elohim, is any one who doeth these things. (Deu. 22:5)

And, with mankind, shalt thou not lie, as with womankind,—an abomination, it is. And, of no beast, shalt thou have carnal knowledge, to commit uncleanness therewith,—neither shall, a woman, present herself to a beast to couch down thereto,—confusion, it is. (Lev. 18:22-23)

⁴ **I. Improving Classroom Instruction** Inclusion of GLBT (*Gay, Lesbian, Bisexual, & Transgender*) -related topics and themes in classroom instruction is an important step in fostering a safe environment for GLBT and gender nonconforming students. One study found that GLBT students' feeling of safety increased when there was explicit mention of GLBT issues in the curriculum. (UC Davis 2004). Inclusion of GLBT-related curricula also helps heterosexual youth to better understand and develop healthy attitudes toward themselves and their fellow students. http://www.nea.org/assets/docs/mf_glbguide.pdf

⁵ As a medical diagnosis, **pedophilia** (or **paedophilia**) is typically defined as a psychiatric disorder in adults (persons age 16 and older) characterized by a primary or exclusive sexual interest in prepubescent children (generally age 13 years or younger, though onset of puberty may vary).

⁶ Homosexual Behavior & Pedophilia; <http://www.us2000.org/cfmc/Pedophilia.pdf>

⁷ OED

⁸ Ge 13:13 Now, the men of Sodom, were base and sinful,—against Yahweh, exceedingly.

And, any man who lieth with mankind as with womankind, an abomination, have both of them wrought,—they shall, surely be put to death,—their blood, shall be upon themselves...And, any man, who hath carnal knowledge of a beast, shall, surely be put to death,—and, the beast, shall ye slay. And, a woman, who approacheth unto any beast, to couch down thereto, then shalt thou slay the woman and the beast,—they shall, surely be put to death,—their blood, shall be upon themselves. (Lev. 20:13-16)

Christ had Paul write about homosexuality by saying, "...for, even their females, exchanged away the natural use into that which is against nature,— In like manner also, even the males, leaving the natural use of the female, flamed out in their eager desire one for another, males with males, the indecency, effecting,—and, the necessary recompence of their error, within themselves, duly receiving... Filled with all unrighteousness, wickedness, greed, baseness, full of envy, murder, strife, deceit, evil disposition, Whisperers, detractors, haters of Yahweh, insolent, arrogant, vain boasters, inventors of vices, unto parents unyielding, Without discernment, regardless of covenants, without natural affection, unmerciful:— Who, indeed, having acknowledged the righteous sentence of Yahweh,—that, they who such things as these do practise, are worthy of death, not only, the same things, are doing, but are even delighting together with them who are practising *them*." (Rm. 1:26-32).

It is indisputable that Yahweh condemns homosexuality and other sexual immoralities, while many in the world promote these sins. Yahweh says, "Flee⁹ porneia¹⁰ [sexual immorality]! Every sin, whatsoever a man shall commit, is, outside his body, but, he that committeth porneia [sexual immorality], bringeth sin into his own body. Or know ye not that, your body, is, a shrine of the holy ruah [spirit] that is within you, which ye have from Yahweh? And ye are not your own; —For ye have been bought with a price! Therefore glorify Yahweh in your body" (1 Cor. 6:18-20). If someone has committed sexual immorality, such as homosexuality, then they can confess this sin to Yahweh and he will forgive them.¹¹

Yahweh has commanded us not to commit such sins, in addition he has also commanded us to remove ourselves from people who commit such sinful acts.¹² This would include not patronizing individuals who are homosexuals by watching television programs, listening to their music, voting for them etc. etc., for we have been called to live holy lives.¹³ We have

⁹ 5343 φευγω pheugo *fyoo'-go* 1) to flee away, seek safety by flight

¹⁰ 4202 πορνεία porneia *por-ni'-ah* 1a) adultery, fornication, homosexuality, lesbianism, intercourse with animals etc.

¹¹ If we are confessing our sins, faithful, is he and, righteous—that he should forgive us our sins, and cleanse us from all unrighteousness. (1 Jn. 1:9)

¹² But, now, I have written unto you not to be mixing yourselves up,—if anyone named a brother, be a fornicator, or covetous, or an idolater, or a reviler, or a drunkard, or an extortioner, with such a one as this, not so much, as to be eating together, (1 Cor. 5:11)

¹³ 1Pe 1:16 Inasmuch as it is written—Holy shall ye be, because, I, am holy.

been called to walk as children of light, reproofing the unfruitful works of darkness, such as homosexuality.¹⁴

As Christians, we must stand firmly against the acceptance of homosexuality in our societies. We must recognize it as an assault from the deceiver to destroy our children. Homosexuality not only separates someone from the light of Yahweh and places them into the darkness of our adversary but it also destroys their very body and the bodies of others, regardless of what mediations science will manufacture in the future. Together, we can shine the light of truth on homosexuality in order that children, women and men will be delivered from this wickedness.

Other sources of information on homosexuality:

The medical information on homosexuality will be presented in Appendix A.

"Homosexual Behavior & Pedophilia" <http://www.us2000.org/cfmc/Pedophilia.pdf>

HIV Surveillance Report: Diagnoses of HIV infection and AIDS in the United States and Dependent Areas, 2009. <http://www.cdc.gov/hiv/topics/surveillance/basic.htm>

HIV among Gay, Bisexual and Other Men Who Have Sex with Men
<http://www.cdc.gov/hiv/topics/msm/pdf/msm.pdf>

¹⁴ For ye were, at one time, darkness, but, now, are ye light in the Lord,—as children of light, be walking, For, the fruit of the light, is in all goodness, and righteousness, and truth,—Putting to the test what is well-pleasing unto the Lord; And be not joining in fellowship in the unfruitful works of darkness, but, rather, be even administering reproof; For, of the secret things which are brought to pass by them, it is, shameful, even to speak! All things, however, when reproofed by the light, become manifest, for, all that of itself maketh manifest, is, light; (Eph. 5:8-13)

Appendix A

The Health Risks of Gay Sex

JOHN R. DIGGS, JR., M.D.

As a physician, it is my duty to assess behaviors for their impact on health and wellbeing. When something is beneficial, such as exercise, good nutrition, or adequate sleep, it is my duty to recommend it. Likewise, when something is harmful, such as smoking, overeating, alcohol or drug abuse, and homosexual sex, it is my duty to discourage it.

Executive Summary

Sexual relationships between members of the same sex expose gays, lesbians and bisexuals to extreme risks of Sexually Transmitted Diseases (STDs), physical injuries, mental disorders and even a shortened life span. There are five major distinctions between gay and heterosexual relationships, with specific medical consequences. They are:

- **Levels of Promiscuity**

Prior to the AIDS epidemic, a 1978 study found that 75 percent of white, gay males claimed to have had more than 100 lifetime male sex partners: 15 percent claimed 100- 249 sex partners; 17 percent claimed 250-499; 15 percent claimed 500- 999; and 28 percent claimed more than 1,000 lifetime male sex partners. Levels of promiscuity subsequently declined, but some observers are concerned that promiscuity is again approaching the levels of the 1970s. The medical consequence of this promiscuity is that gays have a greatly increased likelihood of contracting HIV/AIDS, syphilis and other STDs.

Similar extremes of promiscuity have not been documented among lesbians. However, an Australian study found that 93 percent of lesbians reported having had sex with men, and lesbians were 4.5 times more likely than heterosexual women to have had more than 50 lifetime male sex partners. Any degree of sexual promiscuity carries the risk of contracting STDs.

- **Physical Health**

Common sexual practices among gay men lead to numerous STDs and physical injuries, some of which are virtually unknown in the heterosexual population. Lesbians are also at higher risk for STDs. In addition to diseases that may be transmitted during lesbian sex, a study at an Australian STD clinic found that lesbians were three to four times more likely than heterosexual women to have sex with men who were high-risk for HIV.

- **Mental Health**

It is well established that there are high rates of psychiatric illnesses, including depression, drug abuse, and suicide attempts, among gays and lesbians. This is true even in the Netherlands, where gay, lesbian and bisexual (GLB) relationships are far more socially acceptable than in the U.S. Depression and drug abuse are strongly associated with risky sexual practices that lead to serious medical problems.

- **Life Span**

The only epidemiological study to date on the life span of gay men concluded that gay and bisexual men lose up to 20 years of life expectancy.

- **Monogamy**

Monogamy, meaning long-term sexual fidelity, is rare in GLB relationships, particularly among gay men. One study reported that 66 percent of gay couples reported sex outside the relationship within the first year, and nearly 90 percent if the relationship lasted five years.

Encouraging people to engage in risky sexual behavior undermines good health and can result in a shortened life span. Yet that is exactly what employers and governmental entities are doing when they grant GLB couples benefits or status that make GLB relationships appear more socially acceptable.

The Health Risks of Gay Sex

Introduction

Back in the early 1980s, while working at Beth Israel Hospital, I vividly remember seeing healthy young gay men dying of a mysterious disease that researchers only later identified as a sexually transmitted disease — AIDS. Over the years, I've seen many patients with that diagnosis die.

As a physician, it is my duty to assess behaviors for their impact on health and wellbeing. When something is beneficial, such as exercise, good nutrition, or adequate sleep, it is my duty to recommend it. Likewise, when something is harmful, such as smoking, overeating, alcohol or drug abuse, it is my duty to discourage it.

When sexual activity is practiced outside of marriage, the consequences can be quite serious. Without question, sexual promiscuity frequently spreads diseases, from trivial to serious to deadly. In fact, the Centers for Disease Control and Prevention estimates that 65 million Americans have an incurable sexually transmitted disease (STD).¹

There are differences between men and women in the consequences of same-sex activity. But most importantly, the consequences of homosexual activity are distinct from the consequences of heterosexual activity. As a physician, it is my duty to inform patients of the health risks of gay sex, and to discourage them from indulging in harmful behavior.

I. DIFFERENCES BETWEEN HOMOSEXUAL AND HETEROSEXUAL RELATIONSHIPS

The current media portrayal of gay and lesbian relationships is that they are as healthy, stable and loving as heterosexual marriages — or even more so.² Medical associations are promoting somewhat similar messages.³ Nevertheless, there are at least five major areas of differences between gay and heterosexual relationships, each with specific medical consequences. Those differences include:

- a. Levels of promiscuity
- b. Physical health
- c. Mental health
- d. Life span
- e. Definition of "monogamy"

A. Promiscuity

Gay author Gabriel Rotello notes the perspective of many gays that "Gay liberation was founded . . . on a 'sexual brotherhood of promiscuity,' and any abandonment of that promiscuity would amount to a 'communal betrayal of gargantuan proportions.'"⁴ Rotello's perception of gay promiscuity, which he criticizes, is consistent with survey results. A far-ranging study of homosexual men published in 1978 revealed that 75 percent of self-identified, white, gay men admitted to having sex with more than 100 different males in their lifetime: 15 percent claimed 100-249 sex partners; 17 percent claimed 250- 499; 15 percent claimed 500-999; and 28 percent claimed more than 1,000 lifetime male sex partners.⁵ By 1984, after the AIDS epidemic had taken hold, homosexual men were reportedly curtailing promiscuity, but not by much. Instead of more than 6 partners per month in 1982, the average non-monogamous respondent in San Francisco reported having about 4 partners per month in 1984.⁶

In more recent years, the U.S. Centers for Disease Control has reported an upswing in promiscuity, at least among young homosexual men in San Francisco. From 1994 to 1997, the percentage of homosexual men reporting multiple partners and unprotected anal sex rose from 23.6 percent to 33.3 percent, with the largest increase among men under 25.⁷ Despite its continuing incurability, AIDS no longer seems to deter individuals from engaging in promiscuous gay sex.⁸

The data relating to gay promiscuity were obtained from self-identified gay men. Some advocates argue that the average would be lower if closeted homosexuals were included in the statistics.⁹ That is likely true, according to data obtained in a 2000 survey in Australia that tracked whether men who had sex with men were associated with the gay community. Men who were associated with the gay community were nearly four times as likely to have had more than 50 sex partners in the six months preceding the survey as men who were not associated with the gay community.¹⁰ This may imply that it is riskier to be "out" than "closeted." Adopting a gay identity may create more pressure to be promiscuous and to be so with a cohort of other more promiscuous partners.

Excessive sexual promiscuity results in serious medical consequences — indeed, it is a recipe for transmitting disease and generating an epidemic.¹¹ The HIV/AIDS epidemic has remained a predominantly gay issue in the U.S. primarily because of the greater degree of promiscuity among gays.¹² A study based upon statistics from 1986 through 1990 estimated that 20-year-old gay men had a 50 percent chance of becoming HIV positive by age 55.¹³ As of June 2001, nearly 64 percent of men with AIDS were men who have had sex with men.¹⁴ Syphilis is also more common among gay men. The San Francisco Public Health Department recently reported that syphilis among the city's gay and bisexual men was at epidemic levels. According to the San Francisco Chronicle:

"Experts believe syphilis is on the rise among gay and bisexual men because they are engaging in unprotected sex with multiple partners, many of whom they met in anonymous situations such as sex clubs, adult bookstores, meetings through the Internet and in bathhouses. The new data will show that in the 93 cases involving gay and bisexual men this year, the group reported having 1,225 sexual partners."¹⁵

A study done in Baltimore and reported in the Archives of Internal Medicine found that gay men contracted syphilis at three to four times the rate of heterosexuals.¹⁶ Promiscuity is the factor most responsible for the extreme rates of these and other Sexually Transmitted Diseases cited below, many of which result in a shortened life span for men who have sex with men.

Promiscuity among lesbians is less extreme, but it is still higher than among heterosexual women. Overall, women tend to have fewer sex partners than men. But there is a surprising finding about lesbian promiscuity in the literature. Australian investigators reported that lesbian women were 4.5 times more likely to have had more than 50 lifetime male partners than heterosexual women (9 percent of lesbians versus 2 percent of heterosexual women); and 93 percent of women who identified themselves as lesbian reported a history of sex with men.¹⁷ Other studies similarly show that 75-90 percent of women who have sex with women have also had sex with men.¹⁸

B. Physical Health

Unhealthy sexual behaviors occur among both heterosexuals and homosexuals. Yet the medical and social science evidence indicate that homosexual behavior is uniformly unhealthy. Although both male and female homosexual practices lead to increases in Sexually Transmitted Diseases, the practices and diseases are sufficiently different that they merit separate discussion.

1. Male Homosexual Behavior

Men having sex with other men leads to greater health risks than men having sex with women¹⁹ not only because of promiscuity but also because of the nature of sex among men. A British researcher summarizes the danger as follows:

"Male homosexual behaviour is not simply either 'active' or 'passive,' since penile-anal, mouth-penile, and hand-anal sexual contact is usual for both partners, and mouth-anal contact is not infrequent.... Mouth-anal contact is the reason for the relatively high incidence of diseases caused by bowel pathogens in male homosexuals. Trauma may encourage the entry of micro-organisms and thus lead to primary syphilitic lesions occurring in the anogenital area.... In addition to sodomy, trauma may be caused by foreign bodies, including stimulators of various kinds, penile adornments, and prostheses."²⁰

Although the specific activities addressed below may be practiced by heterosexuals at times, homosexual men engage in these activities to a far greater extent.²¹

a. Anal-genital

Anal intercourse is the sine qua non of sex for many gay men .²² Yet human physiology makes it clear that the body was not designed to accommodate this activity. The rectum is significantly different from the vagina with regard to suitability for penetration by a penis. The vagina has natural lubricants and is supported by a network of muscles. It is composed of a mucus membrane with a multi-layer stratified squamous epithelium that allows it to endure friction without damage and to resist the immunological actions caused by semen and sperm. In comparison, the anus is a delicate mechanism of small muscles that comprise an "exit-only" passage. With repeated trauma, friction and stretching, the sphincter loses its tone and its ability to maintain a tight seal. Consequently, anal intercourse leads to leakage of fecal material that can easily become chronic.

The potential for injury is exacerbated by the fact that the intestine has only a single layer of cells separating it from highly vascular tissue, that is, blood. Therefore, any organisms that are introduced into the rectum have a much easier time establishing a foothold for infection than they would in a vagina. The single layer tissue cannot withstand the friction associated with penile penetration, resulting in traumas that expose both participants to blood, organisms in feces, and a mixing of bodily fluids.

Furthermore, ejaculate has components that are immunosuppressive. In the course of ordinary reproductive physiology, this allows the sperm to evade the immune defenses of the female. Rectal insemination of rabbits has shown that sperm impaired the immune defenses of the recipient.²³ Semen may have a similar impact on humans.²⁴

The end result is that the fragility of the anus and rectum, along with the immunosuppressive effect of ejaculate, make anal-genital intercourse a most efficient manner of transmitting HIV and other infections. The list of diseases found with extraordinary frequency among male homosexual practitioners as a result of anal intercourse is alarming:

- Anal Cancer
- Chlamydia trachomatis
- Cryptosporidium
- Giardia lamblia
- Herpes simplex virus
- Human immunodeficiency virus
- Human papilloma virus
- Isospora belli

Microsporidia
Gonorrhea
Viral hepatitis types B & C
Syphilis²⁵

Sexual transmission of some of these diseases is so rare in the exclusively heterosexual population as to be virtually unknown. Others, while found among heterosexual and homosexual practitioners, are clearly predominated by those involved in homosexual activity. Syphilis, for example is found among heterosexual and homosexual practitioners. But in 1999, King County, Washington (Seattle), reported that 85 percent of syphilis cases were among self-identified homosexual practitioners.²⁶ And as noted above, syphilis among homosexual men is now at epidemic levels in San Francisco.²⁷

A 1988 CDC survey identified 21 percent of all Hepatitis B cases as being homosexually transmitted while 18 percent were heterosexually transmitted.²⁸ Since homosexuals comprise such a small percent of the population (only 1-3 percent),²⁹ they have a significantly higher rate of infection than heterosexuals.³⁰

Anal intercourse also puts men at significant risk for anal cancer. Anal cancer is the result of infection with some subtypes of human papilloma virus (HPV), which are known viral carcinogens. Data as of 1989 showed the rates of anal cancer in male homosexual practitioners to be 10 times that of heterosexual males, and growing.³⁰ Thus, the prevalence of anal cancer among gay men is of great concern. For those with AIDS, the rates are doubled.³¹ Other physical problems associated with anal intercourse are:

hemorrhoids
anal fissures
anorectal trauma
retained foreign bodies.³²

b. Oral-anal

There is an extremely high rate of parasitic and other intestinal infections documented among male homosexual practitioners because of oral-anal contact. In fact, there are so many infections that a syndrome called "the Gay Bowel" is described in the medical literature.³³ "Gay bowel syndrome constitutes a group of conditions that occur among persons who practice unprotected anal intercourse, anilingus, or fellatio following anal intercourse."³⁴ Although some women have been diagnosed with some of the gastrointestinal infections associated with "gay bowel," the vast preponderance of patients with these conditions are men who have sex with men.³⁵

"Rimming" is the street name given to oral-anal contact. It is because of this practice that intestinal parasites ordinarily found in the tropics are encountered in the bodies of American gay men. Combined with anal intercourse and other homosexual practices, "rimming" provides a rich opportunity for a variety of infections.

Men who have sex with men account for the lion's share of the increasing number of cases in America of sexually transmitted infections that are not generally spread through sexual contact. These diseases, with consequences that range from severe and even life-threatening to mere annoyances, include Hepatitis A,³⁶ Giardia lamblia, Entamoeba histolytica,³⁷ Epstein-Barr virus,³⁸

Neisseria meningitides,³⁹ Shigellosis, Salmonellosis, Pediculosis, scabies and Campylobacter.⁴⁰ The U.S. Centers for Disease Control (CDC) identified a 1991 outbreak of Hepatitis A in New York City, in which 78 percent of male respondents identified themselves as homosexual or bisexual.⁴¹ While Hepatitis A can

be transmitted by routes other than sexual, a preponderance of Hepatitis A is found in gay men in multiple states .42 Salmonella is rarely associated with sexual activity except among gay men who have oral-anal and oral-genital contact following anal intercourse.⁴³ The most unsettling new discovery is the reported sexual transmission of typhoid. This water-borne disease, well known in the tropics, only infects 400 people each year in the United States, usually as a result of ingestion of contaminated food or water while abroad. But sexual transmission was diagnosed in Ohio in a series of male sex partners of one male who had traveled to Puerto Rico.⁴⁴

In America, Human Herpes Virus 8 (called Herpes Type 8 or HHV-8) is a disease found exclusively among male homosexual practitioners. Researchers have long noted that men who contracted AIDS through homosexual behavior frequently developed a previously rare form of cancer called Kaposi's sarcoma. Men who contract HIV/AIDS through heterosexual sex or intravenous drug use rarely display this cancer. Recent studies confirm that Kaposi's sarcoma results from infection with HHV-8. The New England Journal of Medicine described one cohort in San Francisco where 38 percent of the men who admitted any homosexual contact within the previous five years tested positive for this virus while none of the exclusively heterosexual men tested positive. The study predicted that half of the men with both HIV and HHV-8 would develop the cancer within 10 years .⁴⁵ The medical literature is currently unclear as to the precise types of sexual behavior that transmit HHV-8, but there is a suspicion that it may be transmitted via saliva.⁴⁶

c. Human Waste

Some gay men sexualize human waste, including the medically dangerous practice of coprophilia, which means sexual contact with highly infectious fecal wastes .⁴⁷ This practice exposes the participants to all of the risks of anal-oral contact and many of the risks of anal-genital contact.

d. Fisting

"Fisting" refers to the insertion of a hand or forearm into the rectum, and is far more damaging than anal intercourse. Tears can occur, along with incompetence of the anal sphincter. The result can include infections, inflammation and, consequently, enhanced susceptibility to future STDs. Twenty-two percent of homosexuals in one survey admitted to having participated in this practice.⁴⁸

e. Sadism

The sexualization of pain and cruelty is described as sadism, named for the 18th Century novelist, the Marquis de Sade. His novel *Justine* describes repeated rapes and non-consensual whipping s.⁴⁹ Not all persons who practice sadism engage in the same activities. But a recent advertisement for a sadistic "conference" included a warning that participants might see "intentional infliction of pain [and] cutting of the skin with bleeding" Scheduled workshops included "Vaginal Fisting" (with a demonstration), "Sacred Sexuality and Cutting" with "a demonstration of a cutting with a live subject," "Rough Rope," and a "Body Harness" workshop that was to involve "demonstrating and coaching the tying of erotic body harnesses that involve the genitals, male and female."⁵⁰ A similar event entitled the "Vicious Valentine" occurred near Chicago on Feb. 15-17, 2002.⁵¹ The medical consequences of such activities range from mild to fatal, depending upon the nature of the injuries inflicted .⁵² As many as 37 percent of homosexuals have practiced some form of sadism .⁵³

f. Conclusion

The consequences of homosexual activity have significantly altered the delivery of medical care to the population at-large. With the increased incidence of STD organisms in unexpected places, simple sore throat is no longer so simple. Doctors must now ask probing questions of their patients or risk making a misdiagnosis. The evaluation of a sore throat must now include questions about oral and anal sex. A case of hemorrhoids is no longer just a surgical problem. We must now inquire as to sexual practice and

consider that anal cancer, rectal gonorrhea, or rectal chlamydia may be secreted in what deceptively appears to be "just hemorrhoids."⁵⁴ Moreover, data shows that rectal and throat gonorrhea, for example, are without symptoms in 75 percent of cases.⁵⁵

The impact of the health consequences of gay sex is not confined to homosexual practitioners. Even though nearly 11 million people in America are directly affected by cancer, compared to slightly more than three-quarters of a million with AIDS,⁵⁶ AIDS spending per patient is more than seven times that for cancer.⁵⁷ The inequity for diabetes and heart disease is even more striking.⁵⁸ Consequently, the disproportionate amount of money spent on AIDS detracts from research into cures for diseases that affect more people.

2. Female Homosexual Behavior

Lesbians are also at higher risk for STDs and other health problems than heterosexuals.⁵⁹ However, the health consequences of lesbianism are less well documented than for male homosexuals. This is partly because the devastation of AIDS has caused male homosexual activity to draw the lion's share of medical attention. But it is also because there are fewer lesbians than gay men,⁶⁰ and there is no evidence that lesbians practice the same extremes of same-sex promiscuity as gay men. The lesser amount of medical data does not mean, however, that female homosexual behavior is without recognized pathology. Much of the pathology is associated with heterosexual activity by lesbians.

Among the difficulties in establishing the pathologies associated with lesbianism is the problem of defining who is a lesbian.⁶¹ Study after study documents that the overwhelming majority of self-described lesbians have had sex with men.⁶² Australian researchers at an STD clinic found that only 7 percent of their lesbian sample had never had sexual contact with a male.⁶³

Not only did lesbians commonly have sex with men, but with lots of men. They were 4.5 times as likely as exclusively heterosexual controls to have had more than 50 lifetime male sex partners.⁶⁴ Consequently, the lesbians' median number of male partners was twice that of exclusively heterosexual women.⁶⁵ Lesbians were three to four times more likely than heterosexual women to have sex with men who were high-risk for HIV disease-homosexual, bisexual, or IV drug-abusing men.⁶⁶ The study "demonstrates that WSW [women who have sex with women] are more likely than non- WSW to engage in recognized HIV risk behaviours such as IDU [intravenous drug use], sex work, sex with a bisexual man, and sex with a man who injects drugs, confirming previous reports."⁶⁷

Bacterial vaginosis, Hepatitis B, Hepatitis C, heavy cigarette smoking, alcohol abuse, intravenous drug use, and prostitution were present in much higher proportions among female homosexual practitioners.⁶⁸ Intravenous drug abuse was nearly six times as common in this group.⁶⁹ In one study of women who had sex only with women in the prior 12 months, 30 percent had bacterial vaginosis.⁷⁰ Bacterial vaginosis is associated with higher risk for pelvic inflammatory disease and other sexually transmitted infections.⁷¹

In view of the record of lesbians having sex with many men, including gay men, and the increased incidence of intravenous drug use among lesbians, lesbians are not low risk for disease. Although researchers have only recently begun studying the transmission of STDs among lesbians, diseases such as "crabs," genital warts, chlamydia and herpes have been reported.⁷² Even women who have never had sex with men have been found to have HPV, trichomoniasis and anogenital warts.⁷³

C. Mental Health

1. Psychiatric Illness

Multiple studies have identified high rates of psychiatric illness, including depression, drug abuse and suicide attempts, among self-professed gays and lesbians.⁷⁴ Some proponents of GLB rights have used these findings to conclude that mental illness is induced by other people's unwillingness to accept same-sex attraction and behavior as normal. They point to homophobia, effectively defined as any opposition to or critique of gay sex, as the cause for the higher rates of psychiatric illness, especially among gay youth.⁷⁵ Although homophobia must be considered as a potential cause for the increase in mental health problems, the medical literature suggests other conclusions.

An extensive study in the Netherlands undermines the assumption that homophobia is the cause of increased psychiatric illness among gays and lesbians. The Dutch have been considerably more accepting of same-sex relationships than other Western countries — in fact, same-sex couples now have the legal right to marry in the Netherlands.⁷⁶ So a high rate of psychiatric disease associated with homosexual behavior in the Netherlands means that the psychiatric disease cannot so easily be attributed to social rejection and homophobia.

The Dutch study, published in the *Archives of General Psychiatry*, did indeed find a high rate of psychiatric disease associated with same-sex sex.⁷⁷ Compared to controls who had no homosexual experience in the 12 months prior to the interview, males who had any homosexual contact within that time period were much more likely to experience major depression, bipolar disorder, panic disorder, agoraphobia and obsessive compulsive disorder. Females with any homosexual contact within the previous 12 months were more often diagnosed with major depression, social phobia or alcohol dependence. In fact, those with a history of homosexual contact had higher rates of nearly all psychiatric pathologies measured in the study.⁷⁸ The researchers found "that homosexuality is not only associated with mental health problems during adolescence and early adulthood, as has been suggested, but also in later life."⁷⁹ Researchers actually fear that methodological features of "the study might underestimate the differences between homosexual and heterosexual people."⁸⁰

The Dutch researchers concluded, "this study offers evidence that homosexuality is associated with a higher prevalence of psychiatric disorders. The outcomes are in line with findings from earlier studies in which less rigorous designs have been employed."⁸¹ The researchers offered no opinion as to whether homosexual behavior causes psychiatric disorders, or whether it is the result of psychiatric disorders.

2. Reckless Sexual Behavior

Depression and drug abuse can lead to reckless sexual behavior, even among those who are most likely to understand the deadly risks. In an article that was part of a series on "AIDS at 20," the *New York Times* reported the risks that many gay men take. One night when a gay HIV prevention educator named Seth Watkins got depressed, he met an attractive stranger, had anal intercourse without a condom — and became HIV positive. In spite of his job training, the HIV educator nevertheless employed the psychological defense of "denial" in explaining his own sexual behavior:

"[L]ike an increasing number of gay men in San Francisco and elsewhere, Mr. Watkins sometimes still puts himself and possibly other people at risk. 'I don't like to think about it because I don't want to give anyone H.I.V.,' Mr. Watkins said."⁸²

Another gay man named Vince, who had never before had anal intercourse without a condom, went to a sex club on the spur of the moment when he got depressed, and had unprotected sex:

"I was definitely in a period of depression And there was just something about that particular circumstance and that particular person. I don't know how to describe it. It just appealed to me; it made it seem like it was all right. "83

Some of the men interviewed by the New York Times are deliberately reckless. One fatalistic gay man with HIV makes no apology for putting other men at risk:

"The prospect of going through the rest of your life having to cover yourself up every time you want to get intimate with someone is an awful one.... Now I've got H.I.V. and I don't have to worry about getting it," he said. "There is a part of me that's relieved. I was tired of always having to be careful, of this constant diligence that has to be paid to intimacy when intimacy should be spontaneous. "84

After admitting to almost never using condoms he adds:

"There is no such thing as safe sex.... If people want to use condoms, they can. I didn't go out and purposely get H.I.V. Accidents happen."85

Other reports show similar disregard for the safety of self and others. A 1998 study in Seattle found that 10 percent of HIV-positive men admitted they engaged in unprotected anal sex, and the percentage doubled in 2000.⁸⁶ According to a study of men who attend gay "circuit" parties, ⁸⁷ the danger at such events is even greater. Ten percent of the men surveyed expected to become HIV- positive in their lifetime. Researchers discovered that 17 percent of the circuit party attendees surveyed were already HIV positive.⁸⁸ Two thirds of those attending circuit parties had oral or anal sex, and 28 percent did not use condoms.⁸⁹

In addition, drug use at circuit parties is ubiquitous. Although only 57 percent admit going to circuit parties to use drugs, 95 percent of the survey participants said they used psychoactive drugs at the most recent event they attended.⁹⁰ There was a direct correlation between the number of drugs used during a circuit party weekend and the likelihood of unprotected anal sex.⁹¹ The researchers concluded that in view of their findings, "the likelihood of transmission of HIV and other Sexually Transmitted Diseases among party attendees and secondary partners becomes a real public health concern." ⁹²

Good mental health would dictate foregoing circuit parties and other risky sex. But neither education nor adequate access to health care is a deterrent to such reckless behavior. "Research at the University of New South Wales found well-educated professional men in early middle age —those who experienced the AIDS epidemic of the 1980s — are most likely not to use a condom."93

D. Shortened Life Span

The greater incidence of physical and mental health problems among gays and lesbians has serious consequences for length of life. While many are aware of the death toll from AIDS, there has been little public attention given to the magnitude of the lost years of life.

An epidemiological study from Vancouver, Canada of data tabulated between 1987 and 1992 for AIDS-related deaths reveals that male homosexual or bisexual practitioners lost up to 20 years of life expectancy. The study concluded that if 3 percent of the population studied were gay or bisexual, the probability of a 20-year-old gay or bisexual man living to 65 years was only 32 percent, compared to 78 percent for men in general.⁹⁴ The damaging effects of cigarette smoking pale in comparison -cigarette smokers lose on average about 13.5 years of life expectancy.⁹⁵

The impact on length of life may be even greater than reported in the Canadian study. First, HIV/AIDS is underreported by as much as 15-20 percent, so it is likely that not all AIDS related deaths were accounted for in the study. 96 Second, there are additional major causes of death related to gay sex. For example, suicide rates among a San Francisco cohort were 3.4 times higher than the general U.S. male population in 1987.97 Other potentially fatal ailments such as syphilis, anal cancer, and Hepatitis B and C also affect gay and bisexual men disproportionately. 98

E. "Monogamy"

Monogamy for heterosexual couples means at a minimum sexual fidelity. The most extensive survey of sex in America found that "a vast majority [of heterosexual married couples] are faithful while the marriage is intact."99 The survey further found that 94 percent of married people and 75 percent of cohabiting people had only one partner in the prior year. 100 In contrast, long-term sexual fidelity is rare among GLB couples, particularly among gay males. Even during the coupling period, many gay men do not expect monogamy. A lesbian critic of gay males notes that:

"After a period of optimism about the long range potential of gay men's one-on-one relationships, gay magazines are starting to acknowledge the more relaxed standards operating here, with recent articles celebrating the bigger bang of sex with strangers or proposing 'monogamy without fidelity'-the latest Orwellian formulation to excuse having your cake and eating it too."101

Gay men's sexual practices appear to be consistent with the concept of "monogamy without fidelity." A study of gay men attending circuit parties showed that 46 percent were coupled, that is, they claimed to have a "primary partner." Twenty-seven percent of the men with primary partners "had multiple sex partners (oral or anal) during their most recent circuit party weekend"102 For gay men, sex outside the primary relationship is ubiquitous even during the first year. Gay men reportedly have sex with someone other than their partner in 66 percent of relationships within the first year, rising to approximately 90 percent if the relationship endures over five years. 103 And the average gay or lesbian relationship is short lived. In one study, only 15 percent of gay men and 17.3 percent of lesbians had relationships that lasted more than three years. 104 Thus, the studies reflect very little long-term monogamy in GLB relationships.

II. CULTURAL IMPLICATIONS OF PROMISCUITY

"Don't tear down a fence until you know why it was put up." ~ African proverb

The societal implications of the unrestrained sexual activity described above are devastating. The ideal of sexual activity being limited to marriage, always defined as male-female, has been a fence erected in all civilizations around the globe.105 Throughout history, many people have climbed over the fence, engaging in premarital, extramarital and homosexual sex. Still, the fence stands; the limits are visible to all. Climbing over the fence, metaphorically, has always been recognized as a breach of those limits, even by the breachers themselves. No civilization can retain its vitality for multiple generations after removing the fence.106

But now social activists are saying that there should be no fence, and that to destroy the fence is an act of liberation.107 If the fence is torn down, there is no visible boundary to sexual expression. If gay sex is socially acceptable, what logical reason can there be to deny social acceptance of adultery, polygamy, or pedophilia? The polygamist movement already has support from some of the advocates for GLB rights.108 And some in the psychological profession are floating the idea that maybe pedophilia is not so damaging to children after all.109

Lesbian social critic Camille Paglia observes, "history shows that male homosexuality, which like prostitution flourishes with urbanization and soon becomes predictably ritualized, always tends toward decadence."¹¹⁰ Gay author Gabriel Rotello writes of the changes in homosexual behavior in the last century:

"Most accounts of male-on-male sex from the early decades of this century [20th] cite oral sex, and less often masturbation, as the predominant forms of activity, with the acknowledged homosexual fellating or masturbating his partner. Comparatively fewer accounts refer to anal sex. My own informal survey of older gay men who were sexually active prior to World War II gives credence to the idea that anal sex, especially anal sex with multiple partners, was considerably less common than it later became."¹¹¹

Not only has the practice of anal sex increased, condom use has declined 20 percent and multipartner sex has doubled in the last seven years,¹¹² despite billions of dollars spent on HIV prevention campaigns. "In many cases, the prevention slogans that galvanized gay men in the early years of the epidemic now fall on deaf ears."¹¹³ As should be expected, the health-care costs resulting from gay promiscuity are substantial.¹¹⁴

Social approval of gay sex leads to an increase in such behavior. As early as 1993, Newsweek reported that the growing media presence and social acceptance of homosexual behavior was leading to teenager experimentation to the extent that it was "becoming chic."¹¹⁵ A more recent report stated that "the way gays and lesbians appear in the media may make some people more comfortable acting on homosexual impulses."¹¹⁶ In addition, one of the goals of GLB advocates' quest for domestic partner benefits from employers is to motivate more gays and lesbians "to come out of the closet."¹¹⁷ If, as suggested above, being "out" results in a greater incidence of promiscuity, employer decisions to provide domestic partner benefits may have a negative impact on employee health. Indeed, giving gays and lesbians the social approval they desire may ultimately lead to an early death for employees who otherwise might have restrained their sexual behavior.

Research designed to prove that gays and lesbians are "born that way" has come up empty — there is no scientific evidence that being gay or lesbian is genetically determined.¹¹⁸ Even researcher Dean Hamer, who once hoped he had identified a "gay gene," admits "there is a lot more than just genes going on."¹¹⁹

CONCLUSION

It is clear that there are serious medical consequences to same-sex behavior. Identification with a GLB community appears to lead to an increase in promiscuity, which in turn leads to a myriad of Sexually Transmitted Diseases and even early death. A compassionate response to requests for social approval and recognition of GLB relationships is not to assure gays and lesbians that homosexual relationships are just like heterosexual ones, but to point out the health risks of gay sex and promiscuity. Approving same-sex relationships is detrimental to employers, employees and society in general.

APPENDIX A

Definitional Impediments to Research

Unfortunately, endeavors to assess the actual practices and the health consequences of male and female homosexual behavior are hampered by imprecise definitions. For many, being gay or lesbian or bisexual

is a political identity that does not necessarily correspond to sexual behavior. And investigators find that sexual behavior fluctuates over time:

"[P]eople often change their sexual behavior during their lifetimes, making it impossible to state that a particular set of behaviors defines a person as gay. A man who has sex with men today, for example, might not have done so 10 years ago."¹²⁰

Defining the terms becomes even more difficult when people who identify as gay or lesbian enter heterosexual relationships. Joanne Loulan, a well-known lesbian, has talked openly about her two-year relationship with a man: "I come from this background that sex is an activity, it's not an identity," says Loulan. "It was funny for a while, but then it turned out to be something more connected, more deep. Something more important. And that's when my life started really going topsy turvy." While critics complain that "You can't be a lesbian and be having sex with men," Loulan sees no contradiction in the fact that she "adamantly refuses to call herself a bisexual, to give up the lesbian identity."¹²¹

Several high-profile lesbian media stars that have abandoned lesbianism further illustrate the difficulty in defining homosexuality. An article about the now defunct couple, Anne Heche and Ellen Degeneres, said, "Although the pair never publicly discussed the reason for their breakup, it has been heavily rumored that Heche decided to go back to heterosexuality."¹²² Heche married a man on Sept. 1, 2001.¹²³

As recently as June 2000, pop-music star Sinéad O'Connor said, "I'm a lesbian ... although I haven't been very open about that, and throughout most of my life I've gone out with blokes because I haven't necessarily been terribly comfortable about being a lesbian. But I actually am a lesbian."¹²⁴ Then, shocking the gay world that applauded her "coming out," O'Connor's sexual identity fluctuated again when she withdrew from participating in a lesbian music festival because of her marriage to British Press Association reporter Nick Sommerlad.¹²⁵

Although women get most of the press coverage of fluctuating between same-sex and heterosexual relationships, men can experience similar fluidity. Gay author John Stoltenberg has lived with a lesbian, Andrea Dworkin, since 1974.¹²⁶ And a 2000 survey in Australia found that 19 percent of gay men reported having sex with a woman in the six months prior to the survey.¹²⁷ This fluctuation in sexual "orientation" inhibits the creation of a fixed definition of homosexuality. As one group of researchers stated the problem:

"Does a man who has homosexual sex in prison count as a homosexual? Does a man who left his wife of twenty years for a gay lover count as a homosexual or heterosexual? Do you count the number of years he spent with his wife as compared to his lover? Does the married woman who had sex with her college roommate a decade ago count? Do you assume that one homosexual experience defines someone as gay for all time?"¹²⁸

Despite the difficulty in defining homosexuality, the one thing that is clear is that those who engage in same-sex practices or identify themselves as gay, lesbian or bisexual constitute a very small percentage of the population. The most reliable studies indicate that 1-3 percent of people — and probably less than 2 percent — consider themselves to be gay, lesbian or bisexual, or currently practice same-sex sex.¹²⁹

End notes

1. "Tracking the Hidden Epidemics: Trends in STDs in the United States, 2000," Centers for Disease Control and Prevention (CDC), available at www.cdc.gov.
2. Becky Birtha, "Gay Parents and the Adoption Option," *The Philadelphia Inquirer*, March 04, 2002, www.philly.com/mld/inquirer/news/editorial/2787531.htm; Grant Pick, "Make Room for Daddy — and Poppa," *The Chicago Tribune* Internet Edition, March 24, 2002, www.chicagotribune.com/features/magazine/chi-0203240463mar24.story
3. Ellen C. Perrin, et al., "Technical Report: Coparent or Second-Parent Adoption by Same-Sex Parents," *Pediatrics*, 109(2): 341-344 (2002).
4. Gabriel Rotello, *Sexual Ecology: AIDS and the Destiny of Gay Men*, p. 112, New York: Penguin Group, 1998 (quoting gay writer Michael Lynch).
5. Alan P. Bell and Martin S. Weinberg, *Homosexualities: A Study of Diversity Among Men and Women*, p. 308, Table 7, New York: Simon and Schuster, 1978.
6. Leon McKusick, et al., "Reported Changes in the Sexual Behavior of Men at Risk for AIDS, San Francisco, 1982-84 — the AIDS Behavioral Research Project," *Public Health Reports*, 100(6): 622-629, p. 625, Table 1 (November- December 1985). In 1982 respondents reported an average of 4.7 new partners in the prior month; in 1984, respondents reported an average of 2.5 new partners in the prior month.
7. "Increases in Unsafe Sex and Rectal Gonorrhoea among Men Who Have Sex with Men — San Francisco, California, 1994-1997," *Mortality and Morbidity Weekly Report*, CDC, 48(03): 45-48, p. 45 (January 29, 1999).
8. This was evident by the late 80's and early 90's. Jeffrey A. Kelly, PhD, et al., "Acquired Immunodeficiency Syndrome/ Human Immunodeficiency Virus Risk Behavior Among Gay Men in Small Cities," *Archives of Internal Medicine*, 152: 2293-2297, pp. 2295-2296 (November 1992); Donald R. Hoover, et al., "Estimating the 1978-1990 and Future Spread of Human Immunodeficiency Virus Type 1 in Subgroups of Homosexual Men," *American Journal of Epidemiology*, 134(10): 1190-1205, p. 1203 (1991).
9. A lesbian pastor made this assertion during a question and answer session that followed a presentation the author made on homosexual health risks at the Chatauqua Institute in Western New York, summer 2001.
10. Paul Van de Ven, et al., "Facts & Figures: 2000 Male Out Survey," p. 20 & Table 20, monograph published by National Centre in HIV Social Research Faculty of Arts and Social Sciences, The University of New South Wales, February 2001.
11. Rotello, pp. 43-46.
12. *Ibid.*, pp. 165-172.
13. Hoover, et al., Figure 3.
14. "Basic Statistics," CDC — Division of HIV/AIDS Prevention, June 2001, www.cdc.gov/hiv/stats.htm. (Nearly 8% (50,066) of men with AIDS had sex with men and used intravenous drugs. These men are included in the 64% figure (411,933) of 649,186 men who have been diagnosed with AIDS.)
15. Figures from a study presented at the Infectious Diseases Society of America meeting in San Francisco and reported by Christopher Heredia, "Big spike in cases of syphilis in S.F.: Gay, bisexual men affected most," *San Francisco Chronicle*, October 26, 2001, www.sfgate.com/cgi-bin/article.cgi?file=/chronicle/archive/2001/10/26/MN74893.DTL.
16. Catherine Hutchinson, et al., "Characteristics of Patients with Syphilis Attending Baltimore STD Clinics," *Archives of Internal Medicine*, 151: 511-516, p. 513 (1991).
17. Katherine Fethers, Caron Marks, et al., "Sexually transmitted infections and risk behaviours in women who have sex with women," *Sexually Transmitted Infections*, 76(5): 345-349, p. 347 (October 2000).
18. James Price, et al., "Perceptions of cervical cancer and pap smear screening behavior by Women's Sexual Orientation," *Journal of Community Health*, 21(2): 89-105 (1996); Daron Ferris, et al., "A Neglected Lesbian Health Concern: Cervical Neoplasia," *The Journal of Family Practice*, 43(6): 581-584, p. 581 (December 1996); C. Skinner, J. Stokes, et al., "A Case-Controlled Study of the Sexual Health Needs of Lesbians," *Sexually Transmitted Infections*, 72(4): 277-280, Abstract (1996).
19. The Gay and Lesbian Medical Association (GLMA) recently published a press release entitled "Ten Things Gay Men Should Discuss with Their Health Care Providers" (July 17, 2002), www.glma.org/news/releases/n02071710gaythings.html. The list includes: HIV/AIDS (Safe Sex), Substance Use, Depression/ Anxiety, Hepatitis Immunization, STDs, Prostate/ Testicular/Colon Cancer, Alcohol, Tobacco, Fitness and Anal Papilloma.

20. R. R. Wilcox, "Sexual Behaviour and Sexually Transmitted Disease Patterns in Male Homosexuals," *British Journal of Venereal Diseases*, 57(3): 167-169, 167 (1981).
21. Robert T. Michael, et al., *Sex in America: a Definitive Survey*, pp. 140-141, Table 11, Boston: Little, Brown, and Co., 1994; Rotello, pp. 75-76.
22. Rotello, p. 92.
23. Jon M. Richards, J. Michael Bedford, and Steven S. Witkin, "Rectal Insemination Modifies Immune Responses in Rabbits," *Science*, 27(224): 390-392 (1984).
24. S. S. Witkin and J. Sonnabend, "Immune Responses to Spermatozoa in Homosexual Men," *Fertility and Sterility*, 39(3): 337-342, pp. 340-341 (1983).
25. Anne Rompalo, "Sexually Transmitted Causes of Gastrointestinal Symptoms in Homosexual Men," *Medical Clinics of North America*, 74(6): 1633-1645 (November 1990); "Anal Health for Men and Women," *LGBTHealthChannel*, www.gayhealthchannel.com/analhealth/; "Safer Sex (MSM) for Men who Have Sex with Men," *LGBTHealthChannel*, www.gayhealthchannel.com/stdmsm/.
26. "Resurgent Bacterial Sexually Transmitted Disease Among Men Who Have Sex With Men — King County, Washington, 1997-1999," *Morbidity and Mortality Weekly Report*, CDC, 48(35): 773-777 (September 10, 1999).
27. Heredia, "Big spike in cases of syphilis in S.F.: Gay, bisexual men affected most."
28. "Changing Patterns of Groups at High Risk for Hepatitis B in the United States," *Morbidity and Mortality Weekly Report*, CDC, 37(28): 429-432, p. 437 (July 22, 1988). Hepatitis B and Care viral diseases of the liver.
29. Edward O. Laumann, John H. Gagnon, et al., *The social organization of sexuality: Sexual practices in the United States*, p. 293, Chicago: University of Chicago Press, 1994; Michael, et al., p. 176; David Forman and Clair Chilvers, "Sexual Behavior of Young and Middle-Aged Men in England and Wales," *British Medical Journal*, 298: 1137-1142 (1989); and Gary Remafedi, et al., "Demography of Sexual Orientation in Adolescents," *Pediatrics*, 89: 714-721 (1992). See appendix A.
30. Mads Melbye, Charles Rabkin, et al., "Changing patterns of anal cancer incidence in the United States, 1940-1989," *American Journal of Epidemiology*, 139: 772-780, p. 779, Table 2 (1994).
31. James Goedert, et al., for the AIDS-Cancer Match Study Group, "Spectrum of AIDS-associated malignant disorders," *The Lancet*, 351: 1833-1839, p. 1836 (June 20, 1998).
32. "Anal Health for Men and Women," *LGBTHealthChannel*, www.gayhealthchannel.com/analhealth/; J. E. Barone, et al., "Management of Foreign Bodies and Trauma of the Rectum," *Surgery, Gynecology and Obstetrics*, 156(4): 453-457 (April 1983).
33. Henry Kazal, et al., "The gay bowel syndrome: Clinicopathologic correlation in 260 cases," *Annals of Clinical and Laboratory Science*, 6(2): 184-192 (1976).
34. Glen E. Hastings and Richard Weber, "Use of the term 'Gay Bowel Syndrome,'" reply to a letter to the editor, *American Family Physician*, 49(3): 582 (1994).
35. Ibid.; E. K. Markell, et al., "Intestinal Parasitic Infections in Homosexual Men at a San Francisco Health Fair," *Western Journal of Medicine*, 139(2): 177-178 (August, 1983).
36. "Hepatitis A among Homosexual Men — United States, Canada, and Australia," *Morbidity and Mortality Weekly Report*, CDC, 41(09): 155, 161-164 (March 06, 1992).
37. Rompalo, p. 1640.
38. H. Naher, B. Lenhard, et al., "Detection of Epstein-Barr virus DNA in anal scrapings from HIV-positive homosexual men," *Archives of Dermatological Research*, 287(6): 608- 611, Abstract (1995).
39. B. L. Carlson, N. J. Fiumara, et al., "Isolation of Neisseria meningitidis from anogenital specimens from homosexual men," *Sexually Transmitted Diseases*, 7(2): 71-73 (April 1980).
40. P. Paulet and G. Stoffels, "Maladies anorectales sexuellement transmissibles" ["Sexually-Transmissible Anorectal Diseases"], *Revue Medicale Bruxelles*, 10(8): 327-334, Abstract (October 10, 1989).
41. "Hepatitis A among Homosexual Men — United States, Canada, and Australia," *Morbidity and Mortality Weekly Report*, CDC, 41(09): 155, 161-164 (March 06, 1992).
42. Ibid.
43. C. M. Thorpe and G. T. Keutsch, "Enteric bacterial pathogens: Shigella, Salmonella, Campylobacter," in K. K. Holmes, P. A. Mardh, et al., (Eds.), *Sexually Transmitted Diseases* (3rd edition), p. 549, New York: McGraw-Hill Health Professionals Division, 1999.
44. Tim Bonfield, "Typhoid traced to sex encounters," *Cincinnati Enquirer*, April 26, 2001; Erin

- McClam, "Health Officials Document First Sexual Transmission of Typhoid in U.S.," *Associated Press*, April 25, 2001, www.thebody.com/cdc/news_updates_archive/apr26_01/typhoid.html. A representative of the Foodborne and Diarrheal Diseases Branch, Division of Bacterial and Mycotic Diseases at the CDC in Atlanta, Georgia, confirmed this report and provided a link to the AP story on October 4, 2002.
45. Jeffrey Martin, et al., "Sexual Transmission and the Natural History of Human Herpes Virus 8 Infection," *New England Journal of Medicine*, 338(14): 948-954, p. 952 (1998).
 46. Alexandra M. Levine, "Kaposi's Sarcoma: Far From Gone," paper presented at 5th International AIDS Malignancy Conference, April 23-25, 2001, Bethesda, Maryland, www.medscape.com/viewarticle/420749.
 47. "Paraphilias," *Diagnostic and Statistical Manual of Mental Disorders*, Fourth Edition, Text Revision, p. 576, Washington: American Psychiatric Association, 2000; Karla Jay and Allen Young, *The Gay Report: Lesbians and Gay Men Speak Out About Sexual Experiences and Lifestyles*, pp. 554-555, New York: Summit Books (1979).
 48. Jay and Young, pp. 554-555.
 49. Sade, Marquis de, *Justine or Good Conduct Well Chastised* (1791), New York: Grove Press (1965).
 50. Michigan Rope internet advertisement for "Bondage and Beyond," which was scheduled for February 9-10, 2002, near Detroit, Michigan, www.michiganrope.com/MichiganRopeWorkshop.html. The explicit nature of the advertisement was changed following unexpected publicity, and the hotel where the conference was scheduled ultimately canceled it. Marsha Low, "Hotel Ties Noose Around 2-Day Bondage Meeting," *Detroit Free Press*, January 25, 2002, www.freep.com/news/locoak/nrope25_20020125.htm.
 51. Allyson Smith, "Ramada to host 'Vicious Valentine' Event," *WorldNet Daily*, February 14, 2002, www.worldnetdaily.com/news/article.asp?ARTICLE_ID=26453; "Vicious Valentine 5 Celebrates Mardi Gras, Feb 15-17, 2002," www.leatherquest.com/events/vv2002.htm.
 52. The sadistic rape of 13-year-old Jesse Dirkhising on September 26, 1999, left him dead. See Andrew Sullivan, "The Death of Jesse Dirkhising," *The Pittsburgh Post-Gazette*, April 1, 2001.
 53. Jay and Young, pp. 554-555.
 54. Gay and Lesbian Medical Association, "MSM: Clinician's Guide to Incorporating Sexual Risk Assessment in Routine Visits," www.glma.org/medical/clinical/msm_assessment.html.
 55. S. Bygdeman, "Gonorrhoea in men with homosexual contacts. Serogroups of isolated gonococcal strains related to antibiotic susceptibility, site of infection, and symptoms," *British Journal of Venereal Diseases*, 57(5): 320-324, Abstract (October 1981).
 56. As of January 1, 1999, the National Cancer Institute (NCI) estimated the cancer prevalence in the United States to be 8.9 million. "Estimated US Cancer Prevalence Counts: Who Are Our Cancer Survivors in the US?," *Cancer Control & Population Sciences*, National Cancer Institute, April 2002, www.cancercontrol.cancer.gov/ocs/prevalence. In 1999, the American Cancer Society (ACS) estimated 1,221,800 new cancer cases in the US and an estimated 563,100 cancer related deaths, "Cancer Facts and Figures 1999," p. 4, American Cancer Society, Inc., 1999, www.cancer.org/downloads/STT/F&F99.pdf; in 2000, the ACS estimated 1,220,100 new cancer cases and 552,200 deaths from cancer, "Cancer Facts and Figures 2000," p. 4, American Cancer Society, Inc., 2000, www.cancer.org/downloads/STT/F&F00.pdf; in 2001, the ACS estimated a total number of 1,268,000 new cases of cancer and 553,400 deaths, "Cancer Facts and Figures 2001," p. 5, American Cancer Society, Inc., 2001, www.cancer.org/downloads/STT/F&F2001.pdf. This results in an estimated growth of 2,041,200 new cancer cases over the past three years and an estimated 10,941,200 people with cancer as of January 1, 2002. In 2001 there were 793,025 reported AIDS cases. "Basic Statistics," CDC — Division of HIV/AIDS Prevention, June 2001, www.cdc.gov/hiv/stats.htm.
 57. The federal spending for AIDS research in 2001 was \$2,247,000,000, while the spending for cancer research was not even double that at \$4,376,400,000. "Funding For Research Areas of Interest," National Institute of Health, 2002, www4.od.nih.gov/officeofbudget/FundingResearchAreas.htm.
 58. *Ibid.*; "Fast Stats A to Z: Diabetes," CDC — National Center for Health Statistics, June 04, 2002, www.cdc.gov/nchs/fastats/diabetes.htm; "Fast Stats A to Z: Heart Disease," CDC — National Center for Health Statistics, June 06, 2002, www.cdc.gov/nchs/fastats/heart.htm.
 59. Gay and Lesbian Medical Association Press Release, "Ten Things Lesbians Should Discuss with Their Health Care Providers" (July 17, 2002), www.glma.org/news/releases/n02071710lesbianthings.html. The list includes Breast Cancer, Depression/Anxiety, Gynecological Cancer, Fitness, Substance Use, Tobacco, Alcohol, Domestic Violence,

Osteoporosis and Heart Health.

60. Michael, et al., p. 176 ("about 1.4 percent of women said they thought of themselves as homosexual or bisexual and about 2.8% of the men identified themselves in this way").
61. See Appendix A.
62. Skinner, et al., Abstract; Ferris, et al. p. 581; James Price, et al., p. 90; see Appendix A.
63. Katherine Fethers, et al., "Sexually transmitted infections and risk behaviours in women who have sex with women," *Sexually Transmitted Infections*, 76(5): 345-349, p. 348 (2000).
64. *Ibid.*, p. 347.
65. *Ibid.*
66. *Ibid.*
67. *Ibid.*, p. 348.
68. *Ibid.*, p. 347, Table 1; Susan D. Cochran, et al., "Cancer- Related Risk Indicators and Preventive Screening Behaviors Among Lesbians and Bisexual Women," *American Journal of Public Health*, 91(4): 591-597 (April 2001); Juliet Richters, Sara Lubowitz, et al., "HIV risks among women in contact with Sydney's gay and lesbian community," *Venereology*, 11(3): 35-38 (1998); Juliet Richters, Sarah Bergin, et al., "Women in Contact with the Gay and Lesbian Community: Sydney Women and Sexual Health Survey 1996 and 1998," National Centre in HIV Social Research, University of New South Wales, 1999.
69. Fethers, et al., p. 347 and Table 1.
70. Barbara Berger, Shelley Kolton, et al., "Bacterial vaginosis in lesbians: a sexually transmitted disease," *Clinical Infectious Diseases*, 21: 1402-1405 (1995).
71. E. H. Koumans, et al., "Preventing adverse sequelae of Bacterial Vaginosis: a Public Health Program and Research Agenda," *Sexually Transmitted Diseases*, 28(5): 292-297 (May 2001); R. L. Sweet, "Gynecologic Conditions and Bacterial Vaginosis: Implications for the Non-Pregnant Patient," *Infectious Diseases in Obstetrics and Gynecology*, 8(3): 184-190 (2000).
72. Kathleen M. Morrow, Ph.D., et al., "Sexual Risk in Lesbians and Bisexual Women," *Journal of the Gay and Lesbian Medical Association*, 4(4): 159-165, p. 161 (2000).
73. *Ibid.*, p. 159.
74. For example, Judith Bradford, Caitlin Ryan, and Esther D. Rothblum, "National Lesbian Health Care Survey: Implications for Mental Health Care," *Journal of Consulting and Clinical Psychology*, 62(2): 228-242 (1994); Richard C. Pillard, "Sexual orientation and mental disorder," *Psychiatric Annals*, 18(1): 52-56 (1988); see also Mubarak S. Dahir, "The Gay Community's New Epidemic," *Daily News* (June 5, 2000), www.gaywired.com/story/detail.cfm?Section=12&ID=148&ShowDate=1.
75. Katherine A. O'Hanlan, M.D., et al., "Homophobia As a Health Hazard," *Report of the Gay & Lesbian Medical Association*, pp. 3, 5, www.ohanlan.com/phobiahzd.htm; Laura Dean, et al., "Lesbian, Gay, Bisexual, and Transgender Health: Findings & Concerns," *Journal of the Gay & Lesbian Medical Association*, 4(3): 102-151, pp. 102, 116 (2000).
76. "Netherlands Ends Discrimination in Civil Marriage: Gays to Wed," Lambda Legal Defense and Education Fund Press Release, March 30, 2001, <http://lambdalegal.org/cgibin/pages/documents/record?record=814>.
77. Theo Sandfort, Ron de Graaf, et al., "Same-sex Sexual Behavior and Psychiatric Disorders," *Archives of General Psychiatry*, 58(1): 85-91, p. 89 and Table 2 (January 2001).
78. *Ibid.*
79. *Ibid.*, p. 89.
80. *Ibid.*, p. 90 (emphasis added).
81. *Ibid.*
82. Erica Goode, "With Fears Fading, More Gays Spurn Old Preventive Message," *New York Times*, August 19, 2001.
83. *Ibid.*
84. *Ibid.*
85. *Ibid.*
86. "Officials Voice Alarm Over Halt in AIDS Decline," *New York Times*, August 14, 2001.
87. "A uniform definition of a circuit party does not exist, partly because such parties continue to evolve. However, a circuit party tends to be a multi-event weekend that occurs each year at around the same time and in the same town" Gordon Mansergh, Grant Colfax, et al., "The Circuit Party Men's Health Survey: Findings and Implications for Gay and Bisexual Men," *American Journal of Public Health*, 91(6): 953-958, p. 953 (June 2001).
88. *Ibid.*, p. 955.
89. *Ibid.*, p. 956.
90. *Ibid.*, pp. 956-957, Tables 2 & 3.

91. *Ibid.*, pp. 956-957.
92. *Ibid.*, p. 957. The authors' recommendation was more education.
93. Julie Robotham, "Safe sex by arrangement as gay men reject condoms," *Sydney Morning Herald*, June 7, 2001. Data source: 2000 Male Out Survey, National Centre in HIV Social Research, Australia.
94. R. S. Hogg, S. A. Strathdee, et al., "Modeling the Impact of HIV Disease on Mortality in Gay and Bisexual Men," *International Journal of Epidemiology*, 26(3): 657-661, p. 659 (1997). Death as the result of HIV infection has dropped significantly since 1996. "Life Expectancy Hits New High in 2000; Mortality Declines for Several Leading Causes of Death," CDC News Release, October 10, 2001, www.cdc.gov/nchs/releases/01news/mort2k.htm. Nevertheless, it remains a significant factor in shortened life expectancy for homosexual practitioners.
95. Press Release, *Smoking costs nation \$150 billion each year in health costs, lost productivity*, CDC, Office of Communication, April 12, 2002, www.cdc.gov/od/oc/media/pressrel/r020412.htm.
96. Hogg, et al., p. 660.
97. *Ibid.*
98. "Hepatitis A vaccination of men who have sex with men — Atlanta, Georgia, 1996-1997," *Morbidity and Mortality Report*, CDC, 47(34): 708-711 (September 4, 1998).
99. Robert T. Michael, et al., p. 89.
100. *Ibid.*, p. 101.
101. Camille Paglia, "I'll take religion over gay culture," *Salon.com* online magazine, June 1998, www.frontpagemag.com/archives/guest_column/paglia/gayculture.htm.
102. Gordon Mansergh, Grant Colfax, et al., p. 955.
103. Joseph Harry, *Gay Couples*, p. 116, New York: Praeger Books, 1984.
104. Marcel T. Sagar, M.D. and Eli Robins, M.D., *Male and Female Homosexuality: A Comprehensive Investigation*, p. 57 Table 4.13, p. 225 Table 12.10, Baltimore: The Williams & Wilkins Company, 1973.
105. The existence of limited homosexual relationships in primitive cultures, or even extensive homosexuality in declining civilizations, such as those cited by advocates of same-sex marriage, does not challenge the existence of a prevailing norm. See, for example, William N. Eskridge, Jr., *The Case for Same-Sex Marriage*, Chapter 2, New York: The Free Press, 1996.
106. Joseph D. Unwin, "Sexual Regulations and Cultural Behaviour," pp. 18-19, reprint of Oxford University Press publication of speech given before the Medical Section of the British Psychological Society, March 27, 1935.
107. For example, see the website of the National Coalition for Sexual Freedom, Inc., www.ncsfreedom.org.
108. "The ACLU believes that criminal and civil laws prohibiting or penalizing the practice of plural marriage violate constitutional protections" 1992 Policy Guide of the ACLU, Policy #91, p. 175.
109. Judith Levine, *Harmful to Minors: The Perils of Protecting Children from Sex*, Minneapolis: University of Minnesota Press, 2002; Bruce Rind, Philip Tromovitch, and Robert Bauserman, "A Meta-Analytic Examination of Assumed Properties of Child Sexual Abuse Using College Samples," *Psychological Bulletin*, 124(1): 22-53 (July 1998).
110. Paglia, June 23, 1998.
111. Rotello, p. 42.
112. Goode, August 19, 2001.
113. *Ibid.*
114. See Michael Hamrick, *The Hidden Costs of Domestic Partner Benefits*, pp. 3-4 (Corporate Resource Council, 2002).
115. David Gelman, et al., "Tune In, Come Out," *Newsweek*, p. 70, November 8, 1993.
116. "Iowa study suggests tolerance of homosexuals is growing," *Associated Press*, March 23, 2001.
117. Sally Kohn, *The Domestic Partnership Organizing Manual for Employee Benefits*, p. 1, the Policy Institute of the National Gay and Lesbian Task Force, www.nglft.org/downloads/dp/dp_99.pdf.
118. John Horgan, "Gay Genes, Revisited," *Scientific American*, p. 26, November 1995.
119. Matthew Brellis, "The Fading 'Gay Gene,'" *The Boston Globe*, March 20, 2002, p. C1.
120. Michael, et al., p. 172.
121. Lynn Scherr, "Lesbian Leader Loves a Man," *ABCNews.com*, April 17, 1998.
122. "Former Lesbian Anne Heche Engaged to Cameraman," *ABCNews.com*, June 1, 2001 (emphasis added), reprinted at www.gaywired.com/index.cfm?linkPage=/storydetail.cfm&Section=68&ID=5304.
123. "The Facts: Anne Heche," *Eonline.msn*, April 1, 2002, www.eonline.com/Facts/People/Bio/0,128,31319,00.html.

124. "Sinead O'Connor to Marry a Man," *Reuters*, June 27, 2000, www.g.co.za/2001.2001.06.27-sinead.html.
125. "Sinead Drops out of Wotapalava Tour," *JAM! Music*, May 31, 2001, www.canoe.ca/JamMusicArtistsO/oconnor_sinead.html.
126. John Stoltenberg, "Living with Andrea Dworkin," *Lambda Book Report*, May/June 1994, reprinted at www.nostatusquo.com/ACLU/dworkin/LivingWithAndrea.html.
127. Julie Robotham, "Safe sex by arrangement as gay men reject condoms," *The Sydney Morning Herald*, June 7, 2001. Data source: "2000 Male Out Survey," National Centre in HIV Social Research, Australia.
128. Michael, et al., p. 172.
129. Edward O. Laumann, John H. Gagnon, et al., *The social organization of sexuality: Sexual practices in the United States*, p. 293, Chicago: University of Chicago Press, 1994; Michael, et al., p. 176; David Forman and Clair Chilvers, "Sexual Behavior of Young and Middle-Aged Men in England and Wales," *British Medical Journal*, 298: 1137-1142 (1989); and Gary Remafedi, et al., "Demography of Sexual Orientation in Adolescents," *Pediatrics*, 89: 714-721(1992).

For additional information about how corporate policies can improve employees' health as well as their work-life balance, please contact Paul Weber at the Corporate Resource Council, (480) 444- 0030.

ACKNOWLEDGEMENT

Dr. John R. Diggs, Jr., "The Health Risks of Gay Sex." Corporate Resource Council (2002).

All rights reserved. Permission is granted to reprint this document in its entirety, with proper attribution.

THE AUTHOR

Dr. John R. Diggs, Jr. is a practicing Internist with first-hand experience in treating many of the problems described in this paper. He also travels and lectures on a variety of medical topics to audiences around the world.

Copyright © 2002 Corporate Resource Council.

AIDS Symptoms, Opportunistic Infections, and Cancer¹⁵

Once the CD4 white blood cell count drops below 200 per cubic milligram of blood, HIV has progressed into full-blown AIDS. At this stage of HIV infection, the immune system cannot adequately defend against infection and certain cancers.

Most AIDS symptoms result from opportunistic infections (OI) that only cause disease when the body's immune system has been significantly weakened. The human body is typically host to a wide range of bacteria, viruses, protozoa and fungi. While many of these are not a threat to a healthy body, they are quick to take advantage of a suppressed immune system.

Common OI causes of AIDS symptoms include bacteria-causing pneumonia, severe thrush, toxoplasmosis, and tuberculosis. In all, symptoms of over 25 clinical conditions can appear with AIDS. Symptoms of opportunistic infections can include:

- chronic cough
- coma
- confusion
- difficulty or pain when swallowing
- loss of coordination
- memory problems
- nausea and vomiting
- seizures
- severe fatigue
- severe or chronic diarrhea
- severe weight loss
- shortness of breath
- vision loss or blindness.

In addition to AIDS and symptoms of opportunistic infections, people with AIDS are vulnerable to cancerous cell growth, especially cancers thought to have a viral origin, such as Kaposi's sarcoma and cervical cancer. AIDS and symptoms of lymphoma (cancer of the lymphatic system) are often comorbid, meaning that they coexist.

When present as AIDS symptoms, cancers are generally extremely aggressive. Kaposi's sarcoma, although rare in the general population, is common in AIDS patients. It manifests itself as round red, brown, or purple spots on the skin or in the mouth in light skinned people. In darker skinned individuals, the cancer appears as a pigmented area.

AIDS and Symptoms in Children

Children with AIDS and HIV symptoms often have the same symptoms as adults. In some cases, however, AIDS symptoms may be most apparent in a failure to thrive or grow. Children with AIDS symptoms will experience the same childhood illnesses as healthy children, but the diseases tend to have more severe symptoms and longer recovery times. Severe and frequent ear infections, pink eye (conjunctivitis) and tonsillitis are all common symptoms of HIV/AIDS in children.

¹⁵ Resources:

AIDS.org. (reviewed 19 September 2004). Opportunistic infections. National Institute of Allergy and Infectious Diseases. (updated 4 April 2005). HIV infection and AIDS: An overview. WebMD.com. (nd). Human immunodeficiency virus (HIV) infection. <http://www.tree.com/health/hiv-aids-symptoms.aspx>